

1965-1966 1967-1968 1969-1970 1971-1972 1973-1974 1975-1976 1977-1978 1979-1980 1981-1982 1983-1984 1985-1986 1987-1988 1989-1990 1991-1992 1993-1994 1995-1996 1997-1998 1999-2000 2001-2002 2003-2004 2005-2006 2007-2008 2009-2010 2011-2012 2013-2014 2015-2016 2017-2018 2019-2020 2021-2022 2023-2024 2025-2026 2027-2028 2029-2030 2031-2032 2033-2034 2035-2036 2037-2038 2039-2040 2041-2042 2043-2044 2045-2046 2047-2048 2049-2050 2051-2052 2053-2054 2055-2056 2057-2058 2059-2060 2061-2062 2063-2064 2065-2066 2067-2068 2069-2070 2071-2072 2073-2074 2075-2076 2077-2078 2079-2080 2081-2082 2083-2084 2085-2086 2087-2088 2089-2090 2091-2092 2093-2094 2095-2096 2097-2098 2099-2100 2101-2102 2103-2104 2105-2106 2107-2108 2109-2110 2111-2112 2113-2114 2115-2116 2117-2118 2119-2120 2121-2122 2123-2124 2125-2126 2127-2128 2129-2130 2131-2132 2133-2134 2135-2136 2137-2138 2139-2140 2141-2142 2143-2144 2145-2146 2147-2148 2149-2150 2151-2152 2153-2154 2155-2156 2157-2158 2159-2160 2161-2162 2163-2164 2165-2166 2167-2168 2169-2170 2171-2172 2173-2174 2175-2176 2177-2178 2179-2180 2181-2182 2183-2184 2185-2186 2187-2188 2189-2190 2191-2192 2193-2194 2195-2196 2197-2198 2199-2200 2201-2202 2203-2204 2205-2206 2207-2208 2209-2210 2211-2212 2213-2214 2215-2216 2217-2218 2219-2220 2221-2222 2223-2224 2225-2226 2227-2228 2229-2230 2231-2232 2233-2234 2235-2236 2237-2238 2239-2240 2241-2242 2243-2244 2245-2246 2247-2248 2249-2250 2251-2252 2253-2254 2255-2256 2257-2258 2259-2260 2261-2262 2263-2264 2265-2266 2267-2268 2269-2270 2271-2272 2273-2274 2275-2276 2277-2278 2279-2280 2281-2282 2283-2284 2285-2286 2287-2288 2289-2290 2291-2292 2293-2294 2295-2296 2297-2298 2299-2300 2301-2302 2303-2304 2305-2306 2307-2308 2309-2310 2311-2312 2313-2314 2315-2316 2317-2318 2319-2320 2321-2322 2323-2324 2325-2326 2327-2328 2329-2330 2331-2332 2333-2334 2335-2336 2337-2338 2339-2340 2341-2342 2343-2344 2345-2346 2347-2348 2349-2350 2351-2352 2353-2354 2355-2356 2357-2358 2359-2360 2361-2362 2363-2364 2365-2366 2367-2368 2369-2370 2371-2372 2373-2374 2375-2376 2377-2378 2379-2380 2381-2382 2383-2384 2385-2386 2387-2388 2389-2390 2391-2392 2393-2394 2395-2396 2397-2398 2399-2400 2401-2402 2403-2404 2405-2406 2407-2408 2409-2410 2411-2412 2413-2414 2415-2416 2417-2418 2419-2420 2421-2422 2423-2424 2425-2426 2427-2428 2429-2430 2431-2432 2433-2434 2435-2436 2437-2438 2439-2440 2441-2442 2443-2444 2445-2446 2447-2448 2449-2450 2451-2452 2453-2454 2455-2456 2457-2458 2459-2460 2461-2462 2463-2464 2465-2466 2467-2468 2469-2470 2471-2472 2473-2474 2475-2476 2477-2478 2479-2480 2481-2482 2483-2484 2485-2486 2487-2488 2489-2490 2491-2492 2493-2494 2495-2496 2497-2498 2499-2500 2501-2502 2503-2504 2505-2506 2507-2508 2509-2510 2511-2512 2513-2514 2515-2516 2517-2518 2519-2520 2521-2522 2523-2524 2525-2526 2527-2528 2529-2530 2531-2532 2533-2534 2535-2536 2537-2538 2539-2540 2541-2542 2543-2544 2545-2546 2547-2548 2549-2550 2551-2552 2553-2554 2555-2556 2557-2558 2559-2560 2561-2562 2563-2564 2565-2566 2567-2568 2569-2570 2571-2572 2573-2574 2575-2576 2577-2578 2579-2580 2581-2582 2583-2584 2585-2586 2587-2588 2589-2590 2591-2592 2593-2594 2595-2596 2597-2598 2599-2600 2601-2602 2603-2604 2605-2606 2607-2608 2609-2610 2611-2612 2613-2614 2615-2616 2617-2618 2619-2620 2621-2622 2623-2624 2625-2626 2627-2628 2629-2630 2631-2632 2633-2634 2635-2636 2637-2638 2639-2640 2641-2642 2643-2644 2645-2646 2647-2648 2649-2650 2651-2652 2653-2654 2655-2656 2657-2658 2659-2660 2661-2662 2663-2664 2665-2666 2667-2668 2669-2670 2671-2672 2673-2674 2675-2676 2677-2678 2679-2680 2681-2682 2683-2684 2685-2686 2687-2688 2689-2690 2691-2692 2693-2694 2695-2696 2697-2698 2699-2700 2701-2702 2703-2704 2705-2706 2707-2708 2709-2710 2711-2712 2713-2714 2715-2716 2717-2718 2719-2720 2721-2722 2723-2724 2725-2726 2727-2728 2729-2730 2731-2732 2733-2734 2735-2736 2737-2738 2739-2740 2741-2742 2743-2744 2745-2746 2747-2748 2749-2750 2751-2752 2753-2754 2755-2756 2757-2758 2759-2760 2761-2762 2763-2764 2765-2766 2767-2768 2769-2770 2771-2772 2773-2774 2775-2776 2777-2778 2779-2780 2781-2782 2783

ATTORNEY: Gregory D. Williams  
General Counsel  
NEW ENGLAND BIOLABS, INC.  
32 Tozer Road  
Beverly, MA 01915  
Telephone: (978) 927-5054 X:292  
Facsimile: (978) 927-1705

EXPRESS MAILING LABEL NO.: EE466580584US

## INTEIN-MEDIATED PROTEIN LIGATION OF EXPRESSED PROTEINS

### RELATED APPLICATIONS

5 *See D1* This Application is a Continuation-In-Part of U.S.S.N.  
08/811,492, filed March 5, 1997 now U.S. Patent No.  
5,834,247, issued November 10, 1998, entitled "Modified  
10 Proteins Comprising Controllable Intervening Protein  
Sequences Or Their Elements Methods of Producing Same and  
Methods For Purification Of A Target Protein Comprised By A  
Modified Protein", and of U.S.S.N. 60/102,413, filed September  
30, 1998, entitled "Intein Mediated Peptide Ligation."

### BACKGROUND OF THE INVENTION

15 The present invention relates to methods of intein-  
mediated ligation of proteins. More specifically, the present  
invention relates to intein-mediated ligation of expressed  
20 proteins containing a predetermined N-terminal residue  
and/or a C-terminal thioester generated via use of one or  
more naturally occurring or modified inteins. Preferably, the  
predetermined residue is cysteine.

25 Inteins are the protein equivalent of the self-splicing  
RNA introns (see Perler et al., *Nucleic Acids Res.* 22:1125-  
1127 (1994)), which catalyze their own excision from a  
precursor protein with the concomitant fusion of the flanking  
protein sequences, known as exteins (reviewed in Perler et al.,

*Curr. Opin. Chem. Biol.* 1:292-299 (1997); Perler, F. B. *Cell* 92(1):1-4 (1998); Xu et al., *EMBO J.* 15(19):5146-5153 (1996)).

Studies into the mechanism of intein splicing led to the development of a protein purification system that utilized thiol-induced cleavage of the peptide bond at the N-terminus of the *Sce* VMA intein (Chong et al., *Gene* 192(2):271-281 (1997)). Purification with this intein-mediated system generates a bacterially-expressed protein with a C-terminal thioester (Chong et al., (1997)). In one application, where it is described to isolate a cytotoxic protein, the bacterially expressed protein with the C-terminal thioester is then fused to a chemically-synthesized peptide with an N-terminal cysteine using the chemistry described for "native chemical ligation" (Evans et al., *Protein Sci.* 7:2256-2264 (1998); Muir et al., *Proc. Natl. Acad. Sci. USA* 95:6705-6710 (1998)).

This technique, referred to as "intein-mediated protein ligation" (IPL), represents an important advance in protein semi-synthetic techniques. However, because chemically-synthesized peptides of larger than about 100 residues are difficult to obtain, the general application of IPL is limited by the requirement of a chemically-synthesized peptide as a ligation partner.

IPL technology would be significantly expanded if an expressed protein with a predetermined N-terminus, such as cysteine, could be generated. This would allow the fusion of

one or more expressed proteins from a host cell, such as bacterial, yeast or mammalian cells.

One method of generating an N-terminal cysteine is with the use of proteases. However, proteases have many disadvantages, such as the possibility of multiple protease sites within a protein, as well as the chance of non-specific degradation. Furthermore, following proteolysis, the proteases must be inactivated or purified away from the protein of interest before proceeding with IPL. (Xu, et al., *Proc. Natl. Acad. Sci. USA* 96(2):388-393 (1999) and Erlandson, et al., *Chem. Biol.*, 3:981-991 (1996))

There is, therefore, a need for an improved intein-mediated protein ligation method which overcomes the noted limitations of current IPL methods and which eliminates the need for use of proteases to generate an N-terminal cysteine residue. Such an improved IPL method would have widespread applicability for the ligation of expressed proteins, for example, labeling of extensive portions of a protein for, among other things, NMR analysis.

## SUMMARY OF THE INVENTION

In accordance with the present invention, there is provided a method for the ligation of expressed proteins utilizing one or more inteins which display cleavage at their N- and/or C-termini. In accordance with the present

invention, such inteins may occur either naturally or may be modified to cleave at their N- and/or C-termini. Inteins displaying N- and/or C-terminal cleavage enable the facile isolation of a protein having a C-terminal thioester and a protein having an N-terminal amino acid residue such as cysteine, respectively, for use in the fusion of one or more expressed proteins. Alternatively, the method may be used to generate a single protein having both a C-terminal thioester and a specified N-terminal amino acid residue, such as cysteine, for the creation of cyclic or polymerized proteins. These methods involve the steps of generating at least one C-terminal thioester-tagged first target protein, generating at least one second target protein having a specified N-terminal amino acid residue, for example cysteine, and ligating these proteins. This method may be used where a single protein is expressed, where, for example, the C-terminal thioester end of the protein is fused to the N-terminal end of the same protein. The method may further include chitin-resin purification steps.

In one preferred embodiment the intein from the RIR1 *Methanobacterium thermoautotrophicum* is modified to cleave at either the C-terminus or N-terminus. The modified intein allows for the release of a bacterially expressed protein during a one-column purification, thus eliminating the need proteases entirely. DNA encoding these modified inteins and plasmids containing these modified inteins are also provided by the instant invention.

## BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a diagram depicting both the N-terminal and C-terminal cleavage reactions which comprise intein-mediated protein ligation. The modified *Mth* RIR1 intein was used to purify both MBP with a C-terminal thioester and T4 DNA ligase with an N-terminal cysteine. The *Mth* RIR1 intein for N-terminal cleavage, intein(N), carried the P-1G/N<sup>134</sup>A double mutation. The full length fusion protein consisting of MBP-intein(N)-CBD was separated from cell extract by binding the CBD portion of the fusion protein to a chitin resin. Overnight incubation in the presence of 100 mM 2-mercaptoethanesulfonic acid (MESNA) induced cleavage of the peptide bond prior to the N-terminus of the intein and created a thioester on the C-terminus of MBP. The C-terminal cleavage vector, intein(C), had the P-1G/C<sup>1</sup>A double mutation. The precursor CBD-intein(C)-T4 DNA ligase was isolated from induced *E. coli* cell extract by binding to a chitin resin as described for N-terminal cleavage. Fission of the peptide bond following the C-terminal residue of the intein at a preferred temperature and pH resulted in the production of T4 DNA ligase with an N-terminal cysteine. Ligation occurred when the proteins containing the complementary reactive groups were mixed and concentrated, resulting in a native peptide bond between the two reacting species.

Figure 2A is a gel depicting the purification of a C-terminal thioester-tagged maltose binding protein (MBP) via a thiol-inducible *Mth* RIR1 intein construct pMRB10G (containing the modified intein, R(N), with P<sup>-1</sup>G/N<sup>134</sup>A mutation) and the purification of T4 DNA ligase having an N-terminal cysteine using the vector pBRL-A (containing the modified intein, R(C), with P<sup>-1</sup>G/C<sup>1</sup>A mutation). Lanes 1-3, purification of maltose binding protein (MBP) (M, 43 kDa) with a C-terminal thioester. Lane 1. ER2566 cells transformed with plasmid pMRB10G following Isopropyl β-D-thiogalactopyranoside (IPTG) induction. Lane 2. Cell extract after passage over a chitin resin. Note that the fusion protein, M-R(N)-B, binds to the resin, where B is the chitin binding domain. Lane 3. Fraction 3 of the elution from the chitin resin following overnight incubation at 4°C in the presence of 100 mM MESNA. Lanes 4-6, purification of T4 DNA ligase (L, 56 kDa) with an N-terminal cysteine. Lane 4. IPTG induced ER2566 cells containing plasmid pBRL-A. Lane 5. Cell extract after application to a chitin resin. B-R(C)-L, the fusion protein, binds to the resin. Lane 6. Elution of T4 DNA ligase with an N-terminal cysteine after overnight incubation at room temperature in pH 7 buffer

Figure 2B is a gel depicting ligation of T4 DNA ligase having an N-terminal cysteine to a C-terminal thioester tagged MBP. Lane 1. Thioester-tagged MBP. Lane 2. T4 DNA ligase with an N-terminal cysteine. Lane 3. Ligation reaction

of MBP (0.8 mM) with T4 DNA ligase (0.8 mM), generating M-L, after overnight incubation at 4°C.

Figure 3 is a gel depicting the effect of induction temperature on the cleaving and/or splicing activity of the *Mth* RIR1 intein or *Mth* RIR1 intein mutants. The *Mth* RIR1 intein or mutants thereof, with 5 native N- and C-terminal extein residues were induced at either 15°C or 37°C. The intein was expressed as a fusion protein (M-R-B, 63 kDa) consisting of N-terminal maltose binding protein (M, 43 kDa), the *Mth* RIR1 intein (R, 15 kDa) and at its C-terminus was the chitin binding domain (B, 5 kDa). Lanes 1 and 2. M-R-B with the unmodified *Mth* RIR1 intein. Note the small amount of spliced product (M-B, 48 kDa). Lanes 3 and 4. *Mth* intein with Pro<sup>-1</sup> replaced with Ala, M-R-B(P<sup>-1</sup>A). Both spliced product (M-B) and N-terminal cleavage product (M) are visible. Lanes 5 and 6. Replacement of Pro<sup>-1</sup> with Gly (M-R-B(P<sup>-1</sup>G)) showed some splicing as well as N- and C-terminal cleavage, M and M-R, respectively. Lanes 7 and 8. The Pro<sup>-1</sup> to Gly and Cys<sup>1</sup> to Ser double mutant, M-R-B(P<sup>-1</sup>G/C<sup>1</sup>S), displayed induction temperature dependent C-terminal cleavage (M-R) activity. Lanes 9 and 10. The M-R-B(P<sup>-1</sup>G/N<sup>134</sup>A) mutant possessed only N-terminal cleavage activity producing M. The *Mth* intein or *Mth* intein -CBD fusion is not visible in this Figure.

Figure 4 is a nucleotide sequence (SEQ ID NO:23) comparison of wild type *Mth* RIR1 intein and synthetic *Mth*



RIR1 intein indicating the location of 61 silent base mutations designed to increase expression in *E. coli*. DNA alignment of the wild type *Mth* RIR1 intein (top strand) and the synthetic *Mth* RIR1 intein (bottom strand). To increase expression levels in *E. coli*, 61 silent base changes were made in 49 separate codons when creating the synthetic gene. The first and last codons of the wild type *Mth* RIR1 intein are shown in bold.

### DETAILED DESCRIPTION

The present invention provides a solution to the limitations of current intein-mediated ligation methods by eliminating the need for a synthetic peptide as a ligation partner, and providing a method which is suitable for the fusion one or more expressed proteins.

In general, any intein displaying N- and/or C-terminal cleavage at its splice junctions can be used to generate a defined N-terminus, such as cysteine as well as a C-terminal thioester for use in the fusion of expressed proteins. Inteins which may be used in practicing the present invention include those described in Perler, et al., *Nucleic Acids Res.*, 27(1):346-347 (1999).

In accordance with one preferred embodiment, an intein found in the ribonucleoside diphosphate reductase gene of *Methanobacterium thermoautotrophicum* (the *Mth* RIR1 intein)

was modified for the facile isolation of a protein with an N-terminal cysteine for use in the *in vitro* fusion of two bacterially-expressed proteins. The 134-amino acid *Mth* RIR1 intein is the smallest of the known mini-inteins, and may be close to the minimum amino acid sequence needed to promote splicing (Smith et.al., *J. Bacteriol.* 179: 7135-7155 (1997)).

The *Mth* RIR1 intein has a proline residue on the N-terminal side of the first amino acid of the intein. This residue was previously shown to inhibit splicing in the *Sce* VMA intein (Chong et al., *J. Biol. Chem.* 273:10567-10577 (1998)). The intein was found to splice poorly in *E. coli* when this naturally occurring proline is present. Splicing proficiency increases when this proline is replaced with an alanine residue. Constructs that display efficient N- and C-terminal cleavage are created by replacing either the C-terminal asparagine or N-terminal cysteine of the intein, respectively, with alanine.

These constructs allow for the formation of an intein-generated C-terminal thioester on a first target protein and an intein-generated N-terminal cysteine on a second target protein. These complementary reactive groups may then be ligated via native chemical ligation to produce a peptide bond (Evans et al *supra* (1998), Muir et al *supra* (1998)). Alternatively, a single protein containing both reactive groups may be generated for the creation of cyclic or polymerized

proteins. Likewise, more than one first or second target proteins may be generated via use of multiple mutant inteins.

As used herein, the terms fusion and ligation are used interchangeably. Also as used herein, protein shall mean any protein, fragment of any protein, or peptide capable of ligation according to the methods of the instant invention. Further, as used herein, target protein shall mean any protein the ligation of which, according to the methods of the instant invention, is desired.

The general method of intein-mediated protein ligation in accordance with the present invention is as follows:

(1) An intein of interest is isolated and cloned into an appropriate expression vector(s) such as bacterial, plant, insect, yeast and mammalian cells.

(2) The intein is engineered for N- and/or C-terminal cleavage unless the wild type intein displays the desired cleavage activities. In a preferred embodiment, a modified intein with the desired cleavage properties can be generated by substituting one or more residues within and/or flanking the intein sequence. For example, a modified intein having N-terminal cleavage activity can be created by changing the last intein residue. Alternatively, a modified intein with C-terminal cleavage activity can be created by changing the first intein residue.

(3) The intein with N- and/or C-terminal cleavage activity is fused with an affinity tag to allow purification away from other endogenous proteins.

5

(4) The intein or inteins, either wild type or modified, that display N-terminal and/or C-terminal cleavage, or both, are fused to the desired target protein coding region or regions upstream and/or downstream of the intein.

10

(5) An intein that cleaves at its N-terminus in a thiol reagent dependent manner is used to isolate a protein with a C-terminal thioester. This cleavage and isolation is, for example, carried out as previously described for the *Sce* VMA and *Mxe* GyrA inteins (Chong et al., *Gene* 192(2):271-281 (1997); Evans et al., *Protein Sci.* 7:2256-2264 (1998)). As discussed previously, multiple C-terminal thioester-tagged proteins may be generated at this step .

15

20

(6) A target protein having a specified N-terminus is generated by cleavage of a construct containing an intein that cleaves at its C-terminus. The specified N-terminal residue may be any of the amino acids, but preferably cysteine. As discussed previously, this step may alternately generate a specified N-terminal on the same protein containing a C-terminal thioester, to yield a single protein containing both reactive groups. Alternatively, multiple proteins having the specified N-terminus may be generated at this step.

25

6629" 6456460

(7) Thioester-tagged target protein and target protein having a specified N-termini are fused via intein-mediated protein ligation (IPL) (see Figure 2B). In a preferred embodiment, the N-terminus is cysteine. Alternatively, a single protein containing both a C-terminal thioester and a specified N-terminus, such as a cysteine, may undergo intramolecular ligation to yield a cyclic product and/or intermolecular ligation to yield polymerized proteins.

The methodology described by the instant invention significantly expands the utility of current IPL methods to enable the labeling of extensive portions of a protein for NMR analysis and the isolation of a greater variety of cytotoxic proteins. In addition, this advance opens the possibility of labeling the central portion of a protein by ligating three or more fragments.

The use of an intein or inteins with N-terminal and C-terminal cleavage activity provides the potential to create a defined N-terminus, such as a cysteine, and a C-terminal thioester on a single protein. The intramolecular ligation of the resulting protein generates a circular protein, whereas the intermolecular ligation of several of these proteins generates a protein polymer.

Cleavage at the N- and/or the C-terminus of an intein can be brought about by introducing changes to the intein

and/or its extein sequences. Also, naturally occurring inteins may display these properties and require no manipulation. Cleavage at the N- and/or C-terminus of an intein can occur uncontrollably or induced using nucleophilic compounds, such as thiol reagents, temperature, pH, salt, chaotropic agents, or any combination of the aforementioned conditions and/or reagents.

The Examples presented below are only intended as specific preferred embodiments of the present invention and are not intended to limit the scope of the invention except as provided in the claims herein. The present invention encompasses modifications and variations of the methods taught herein which would be obvious to one of ordinary skill in the art.

The references cited above and below are herein incorporated by reference.

#### EXAMPLE I

##### Creation of the *Mth* RIR1 synthetic gene

The gene encoding the *Mth* RIR1 intein along with 5 native N- and C-extein residues (Smith et al. *supra* (1997)) was constructed using 10 oligonucleotides (New England Biolabs, Beverly, MA) comprising both strands of the gene, as follows:

9) 5'-pCACGTTACAGGTGCGGAAGAAACCTGAGGGGCATGGGTA  
GCCAGAGCCGCGAATCAGTGCGGTGAACGGTTTGCCCTCC  
AGTTCAGCCACAGTGCG-3' (SEQ ID NO:9)

- 10) 5'-pCGGACCGCCACTAGTCATTACAATGGTGTACCGGATACG  
CAGGGGTTGGTTGCC-3' (SEQ ID NO:10)

5 To ensure maximal *E. coli* expression, the coding region  
of the synthetic *Mth* RIR1 intein incorporates 61 silent base  
mutations in 49 of the 134 codons (see Figure 4) in the  
wildtype *Mth* RIR1 intein gene (GenBank AE000845). The  
oligonucleotides were annealed by mixing at equimolar ratios  
10 (400 nM) in a ligation buffer (50 mM Tris-HCl, pH 7.5  
containing 10 mM MgCl<sub>2</sub>, 10 mM dithiothreitol, 1 mM ATP, and  
25 µg BSA) followed by heating to 95°C. After cooling to room  
temperature, the annealed and ligated oligonucleotides were  
inserted into the *Xho*I and *Age*I sites of pMYB5 (NEB), replacing  
the *Sce* VMA intein and creating the plasmid pMRB8P.

15 **Engineering the *Mth* RIR1 intein for N- and C-terminal  
cleavage**

20 The unique *Xho*I and *Spe*I sites flanking the N-terminal  
splice junction and the unique *Bsr*GI and *Age*I sites flanking  
the C-terminal splice junction allowed substitution of amino  
acid residues by linker replacement. The proline residue, Pro-  
1, preceding the intein in pMRB8P was substituted with  
alanine or glycine to yield pMRB8A and pMRB8G1, respectively.  
25 Substitution of Pro-1-Cys<sup>1</sup> with Gly-Ser or Gly-Ala yielded  
pMRB9GS and pMRB9GA, respectively. Replacing Asn<sup>134</sup> with  
Ala in pMRB8G1 resulted in pMRB10G. The following linkers  
were used for substitution of the native amino acids at the



- 16 -

splice junctions (each linker was formed by annealing two synthetic oligonucleotides as described above):

5 P-1A linker: 5'-TCGAGGCAACCAACGCATGCGTATCCGGT  
GACACCATTGTAATGA-3' (SEQ ID NO:11)

and 5'-CTAGTCATTACAATGGTGTCAACCGGATAC  
GCATGCGTTGGTTGCC-3' (SEQ ID NO:12)

10 P-1G linker: 5'-TCGAGGGCTGCGTATCCGGTGACACCATT  
GTAATGA-3 (SEQ ID NO:13)

and 5'-CTAGTCATTACAATGGTGTCAACCGGATAC  
GCAGCCC-3' (SEQ ID NO:14)

15 P-1G/C<sup>1</sup>S linker: 5'-TCGAGGGCATCGAGGCAACCAACGGATC  
CGTATCCGGTGACACCATTGTAATGA-3'  
(SEQ ID NO:15)

20 and 5'-CTAGTCATTACAATGGTGTCAACCGGATAC  
GGATCCGTTGGTTGCCTCGATGCCC-3'  
(SEQ ID NO:16)

25 P-1G/C<sup>1</sup>A linker: 5'-TCGAGGGCATCGAGGCAACCAACGGCGCC  
GTATCCGGTGACACCATTGTAATGA -3'  
(SEQ ID NO:17)

and 5'-CTAGTCATTACAATGGTGTCAACCGGATAC  
GGCGCCGTTGGTTGCCTCGATGCCC-3'  
(SEQ ID NO:18)

30 N<sup>134</sup>A linker: 5'-GTACACGCATGCGGCGAGCAGCCCGG  
GA-3'  
(SEQ ID NO:19)

35

and 5'-CCGGTCCCGGGCTGCTCGCCGCATGC  
GT-3'  
(SEQ ID NO:20)

pBRL-A was constructed by substituting the *Escherichia coli* maltose binding protein (MBP) and the *Bacillus circulans* chitin binding domain (CBD) coding regions in pMRB9GA with the CBD and the T4 DNA ligase coding regions, respectively, subcloned from the pBYT4 plasmid.

## EXAMPLE II

### Generating a thioester-tagged protein:

The pMRB10G construct from Example I contains the *Mth* RIR1 intein engineered to undergo thiol reagent induced cleavage at the N-terminal splice junction (Figure 1, N-terminal cleavage) and was used to isolate proteins with a C-terminal thioester as described previously for the *Sce* VMA and *Mxe* GyrA inteins (Chong et al. *supra* 1997); Evans et al., *supra* (1998)). Briefly, ER2566 cells (Evans et.al. (1998)) containing the appropriate plasmid were grown at 37°C in LB broth containing 100 µg/mL ampicillin to an OD<sub>600</sub> of 0.5-0.6 followed by induction with IPTG (0.5 mM). Induction was either overnight at 15°C or for 3 hours at 30°C.

The cells were pelleted by centrifugation at 3,000xg for 30 minutes followed by resuspension in buffer A (20 mM Tris-HCl, pH 7.5 containing 500 mM NaCl). The cell contents were released by sonication. Cell debris was removed by

centrifugation at 23,000xg for 30 minutes and the supernatant was applied to a column packed with chitin resin (10 mL bed volume) equilibrated in buffer A. Unbound protein was washed from the column with 10 column volumes of buffer A.

5

Thiol reagent-induced cleavage was initiated by rapidly equilibrating the chitin resin in buffer B (20 mM Tris-HCl, pH 8 containing 500 mM NaCl and 100 mM 2-mercaptoethane-sulfonic acid (MESNA)). The cleavage reaction, which simultaneously generates a C-terminal thioester on the target protein, proceeded overnight at 4°C after which the protein was eluted from the column. The use of the pMRB10G construct resulted in the isolation of MBP with a C-terminal thioester (Figure2A).

10

15

### Isolating proteins with an N-terminal cysteine

20

25

The pBRL-A construct from Example I contains an *Mth* RIR1 intein engineered to undergo controllable cleavage at its C-terminus, and was used to purify proteins with an N-terminal cysteine (Figure 1, C-terminal cleavage). The expression and purification protocol was performed as described in Example II, except with buffer A replaced by buffer C (20 mM Tris-HCl, pH 8.5 containing 500 mM NaCl) and buffer B replaced by buffer D (20 mM Tris-HCl, pH 7.0 containing 500 mM NaCl). Also, following equilibration of the

column in buffer D the cleavage reaction proceeded overnight at room temperature.

The expression of plasmid pBRL-A resulted in the purification of 4-6 mg/L cell culture of T4 DNA ligase possessing an N-terminal cysteine (Figure 2A). Protein concentrations were determined using the Bio-Rad protein assay (Bio-Rad Laboratories, Inc., Hercules, CA).

### EXAMPLE III

#### **Protein-protein ligation using Intein-mediated Protein Ligation**

Intein-mediated protein ligation (IPL) was used to fuse two proteins (Figure 2B). Freshly isolated thioester-tagged protein from Example II was mixed with freshly isolated protein containing an N-terminal cysteine residue from Example II, with typical starting concentrations of 1-200  $\mu$ M. The solution was concentrated with a Centriprep 3 or Centriprep 30 apparatus (Millipore Corporation, Bedford, MA) then with a Centricon 3 or Centricon 10 apparatus to a final concentration of 0.15-1.2 mM for each protein.

Ligation reactions proceeded overnight at 4°C and were visualized using SDS-PAGE with 12% Tris-glycine gels (Novex Experimental Technology, San Diego, CA) stained with Coomassie Brilliant Blue. Typical ligation efficiencies ranged from 20-60%.

### Confirmation of ligation in IPL reactions

5 A Factor Xa site in MBP that exists 5 amino acids N-terminal from the site of fusion (Maina et al, *supra* (1988)) allowed amino acid sequencing through the ligation junction. The sequence obtained was NH<sub>2</sub>-TLEGCGEQPTGX<sub>2</sub>LK-COOH (SEQ ID NO:21) which matched the last 4 residues of MBP (TLEG) followed by a linker sequence (CGEQPTG (SEQ ID NO:22)) and the start of T4 DNA ligase (ILK). During amino acid sequencing, the cycle expected to yield an isoleucine did not have a strong enough signal to assign it to a specific residue, so it was represented as an X. The cysteine was identified as the acrylamide alkylation product.

15 The Factor Xa proteolysis was performed on 2 mg of ligation reaction involving MBP and T4 DNA ligase. This reaction mixture was bound to 3 mL of amylose resin (New England Biolabs, Inc., Beverly, MA) equilibrated in buffer A (see Example II). Unreacted T4 DNA ligase was rinsed from the column with 10 column volumes of buffer A. Unligated MBP and the MBP-T4 DNA ligase fusion protein were eluted from the amylose resin using buffer E (20 mM Tris-HCl, pH 7.5 containing 500 mM NaCl and 10 mM maltose). Overnight incubation of the eluted protein with a 200:1 protein:bovine Factor Xa (NEB) ratio (w/w) at 4°C resulted in the proteolysis of the fusion protein and regeneration of a band on SDS-PAGE gels that ran at a molecular weight similar to T4 DNA ligase.

N-terminal amino acid sequencing of the proteolyzed fusion protein was performed on a Procise 494 protein sequencer (PE Applied Biosystems, Foster City, CA).

### Temperature sensitivity of the *Mth* RIR1 intein

The cleavage and/or splicing activity of the *Mth* RIR1 intein was more proficient when protein synthesis was induced at 15°C than when the induction temperature was raised to 37°C (Figure 3). The effect temperature has on the *Mth* RIR1 represents a way to control the activity of this intein for use in controlled splicing or cleavage reactions. Replacement of Pro<sup>-1</sup> with a Gly and Cys<sup>1</sup> with a Ser resulted in a double mutant, the pMRB9GS construct, which showed only *in vivo* C-terminal cleavage activity when protein synthesis was induced at 15°C but not at 37°C. Another double mutant, the pMRB9GA construct, displayed slow cleavage, even at 15°C, which allowed the accumulation of substantial amounts of the precursor protein and showed potential for use as a C-terminal cleavage construct for protein purification.